



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FOSSIL CEPHALOPODA, DESCRIBED BY HYATT
AND CRAGIN, IN THE MUSEUM OF THE
UNIVERSITY OF TEXAS.¹

ANNIE H. PRITCHETT.

In the Second and Third Reports of the Geological Survey of Texas, published in 1890 and 1892 respectively, Professor Alpheus Hyatt described a number of Nautiloids from material collected in Texas under the direction of the then State Geologist, Mr. E. T. Dumble; and in the Fourth Report, Professor F. W. Cragin describes a number of forms, mainly Ammonites. The material described in these three papers represents only a portion, about an eighth, of a rather extensive collection of specimens of these groups, but the portion that is most interesting because of the number of type specimens. Only recently has this material been unpacked and arranged, and because it is the largest collection of the kind in Texas, and because for so many years it has been inaccessible to students, it seemed desirable to publish a short list of the more important specimens. The task has been relatively simple, since in only a few cases had the original labels been lost or misplaced; and in most cases the describers had marked new species with the word "type."

In the following list the name of each species is followed in parentheses by the name of the particular describer, and the volume and page of the published Report.

NAUTILOIDEA.

Temnocheilus conchiferous, type (Hyatt, Second Rep., p. 329).

T. forbesianus McChesney (Hyatt, Second Rep., p. 330).

Tainoceras cavatum, type (Hyatt, Second Rep., p. 341).

T. quadrangulum McChesney (Hyatt, Fourth Rep., p. 402).

Stearoceras gibbosum, type (Hyatt, Fourth Rep., p. 424).

Coloceras globulare, type (Hyatt, Fourth Rep., p. 452).

Glyphioceras cumminsi, type (Hyatt, Fourth Rep., p. 467).

G. incisum, type (Hyatt, Fourth Rep., p. 471).

¹ Contributions from the Zoölogical Laboratory of the University of Texas, No. 66.

- Phacoceras dumbli*, type (Hyatt, Second Rep., p. 347).
Gastrioceras compressum, type (Hyatt, Second Rep., p. 355).
G. entogonum Gabb (Hyatt, Fourth Rep., p. 472).
Metacoceras walcotti, type (Hyatt, Second Rep., p. 337).
M. inconspicuum, type (Hyatt, Second Rep., p. 340).
Domatoceras militarium, type (Hyatt, Fourth Rep., p. 445).
 (The type specimen of *Domatoceras simplex* Hyatt, Fourth Rep., p. 441, appears to have been lost.)
Ephippioceras divisum White and St. John (Hyatt, Second Rep., p. 350).
Solenocheilus collectus Meek and Worthen (Hyatt, Fourth Rep., p. 463).
Paralegoceras iowense Meek and Worthen (Hyatt, Fourth Rep., p. 474).
Nautilus texanus Shumard (Cragin, Fourth Rep., p. 236).

AMMONITOIDEA.

- Placenticeras syrtalis cumminsi*, type (Cragin, Fourth Rep., p. 237).
Pulchella bentonianum, type (Cragin, Fourth Rep., p. 239).
Scaphites septum-seriatus, type (Cragin, Fourth Rep., p. 240).
Sphenodiscus dumbli, type (Cragin, Fourth Rep., p. 243).
S. ræmeri, type (Cragin, Fourth Rep., p. 245).
 (The type specimen of *Sphenodiscus emarginatus*, Cragin, Fourth Rep., p. 245, appears to have been lost.)
S. lenticularis Owen (Cragin, Fourth Rep., p. 245).
Hoplites ræmeri, type (Cragin, Fourth Rep., p. 234).
H. texanus, type (Cragin, Fourth Rep., p. 235).
Pachydiscus brazoensis Shumard (Cragin, Fourth Rep. p. 236).
P. complexus Hall and Meek (Cragin, Fourth Rep., p. 237).
Crioceras annulatus Shumard (Cragin, Fourth Rep., p. 234).
Schlenbachia peruviana Von Buch (Cragin, Fourth Rep., p. 242).
S. woolgari Mantell (Cragin, Fourth Rep., p. 243).
S. leonensis anatina Conrad (Cragin, Fourth Rep., p. 241).
Buchiceras inæquiplacatus Shumard (Cragin, Fourth Rep., p. 233).
B. swallowi Shumard (Cragin, Fourth Rep., p. 234).